

LECTURE 4: DERIVATIVES

Juan Dodyk

WashU

PLAN

Derivatives

1. Definition
2. Calculation
3. Higher Order Derivatives

Optimization

1. Increasing/decreasing
2. Convex/concave
3. Local minima/maxima

L'Hôpital's Rule





TANGENT LINE

Consider the function $f(x) = x^2$.

We want to calculate the slope of the line that “touches” it at $x_0 = 1$.

We can approximate the tangent by lines that pass through $(x_0, f(x_0))$ and $(x, f(x))$ with $x \rightarrow x_0$.

The slope is $\frac{f(x) - f(x_0)}{x - x_0}$.



DERIVATIVES

It makes sense to define the **derivative** of f at x_0 as the limit

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$$

if it exists. If it exists and is not infinite, we say that f is **differentiable** at x_0 .

Note that $x \rightarrow x_0$ is the same as $h \equiv x - x_0 \rightarrow 0$, and $x = x_0 + (x - x_0) = x_0 + h$, so we can write

$$f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}.$$

EXAMPLE

If $f(x) = ax + b$ we have

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{a(x+h) + b - (ax+b)}{h} = \lim_{h \rightarrow 0} \frac{ah}{h} \\ &= \lim_{h \rightarrow 0} a = a. \end{aligned}$$

If $f(x) = x^2$ then

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h} = \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - x^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{2xh + h^2}{h} = \lim_{h \rightarrow 0} (2x + h) = 2x. \end{aligned}$$


DIFFERENTIABLE \Rightarrow CONTINUOUS

We should think of the derivative as the rate of change of $f(x)$ locally.

Notice that if $f'(x_0)$ exists then f has to be continuous at x_0 , because

$$\begin{aligned}\lim_{x \rightarrow x_0} f(x) &= \lim_{x \rightarrow x_0} (f(x) - f(x_0)) + f(x_0) \\ &= \lim_{x \rightarrow x_0} \left[\frac{f(x) - f(x_0)}{x - x_0} (x - x_0) \right] + f(x_0) \\ &= f'(x) \cdot 0 + f(x_0) = f(x_0).\end{aligned}$$

But not every continuous function is differentiable. For example, $f(x) = |x|$ is not differentiable at 0.



CALCULATING DERIVATIVES

If f and g are differentiable at x , $(f \pm g)'(x) = f'(x) \pm g'(x)$. Why?

$$\begin{aligned}(f \pm g)'(x) &= \lim_{h \rightarrow 0} \frac{(f(x+h) \pm g(x+h)) - (f(x) \pm g(x))}{h} \\ &= \lim_{h \rightarrow 0} \left(\frac{f(x+h) - f(x)}{h} \pm \frac{g(x+h) - g(x)}{h} \right) = f'(x) \pm g'(x).\end{aligned}$$

CALCULATING DERIVATIVES

If f and g are differentiable at x , $(fg)'(x) = f'(x)g(x) + f(x)g'(x)$:

$$\begin{aligned}(fg)'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x+h) + f(x)g(x+h) - f(x)g(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{(f(x+h) - f(x))g(x+h) - f(x)(g(x+h) - g(x))}{h} \\&= \lim_{h \rightarrow 0} \left[\frac{f(x+h) - f(x)}{h} g(x+h) + f(x) \frac{g(x+h) - g(x)}{h} \right] \\&= f'(x)g(x) + f(x)g'(x).\end{aligned}$$

Example. $(x^3)' = (x \cdot x^2)' = x' \cdot x^2 + x \cdot (x^2)' = x^2 + x \cdot (2x) = 3x^2$.

In general, $(x^n)' = nx^{n-1}$.

EXERCISE

QUESTION 1

Calculate the derivative of the following functions:

1. $f(t) = 14t - 7$

2. $f(y) = y^3 + 3y^2 - 12$

3. $f(x) = (x^2 + 1)(x^3 - 1)$

4. $f(x) = x^6 + 5x^5 - 2x^2 + 8$

CHAIN RULE

If g is differentiable at x and f is differentiable at $g(x)$, define $(f \circ g)(x) = f(g(x))$, and

$$\begin{aligned}(f \circ g)'(x) &= \lim_{h \rightarrow 0} \frac{f(g(x+h)) - f(g(x))}{h} \\&= \lim_{h \rightarrow 0} \left[\frac{f(g(x+h)) - f(g(x))}{g(x+h) - g(x)} \frac{g(x+h) - g(x)}{h} \right] \\&= f'(g(x))g'(x).\end{aligned}$$

For example, if $f(x) = \frac{1}{x}$ we have

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{1}{x+h} - \frac{1}{x}}{h} = \lim_{h \rightarrow 0} \frac{\frac{x-(x+h)}{(x+h)x}}{h} = \lim_{h \rightarrow 0} \frac{-h}{(x+h)xh} = \lim_{h \rightarrow 0} \frac{-1}{(x+h)x} = -\frac{1}{x^2}.$$

Therefore, $\left(\frac{1}{g(x)}\right)' = (f \circ g)'(x) = f'(g(x))g'(x) = -\frac{1}{g(x)^2}g'(x) = -\frac{g'(x)}{g(x)^2}.$

CALCULATING DERIVATIVES

Notice that if a is constant then $a' = 0$.

Hence if a is constant then $(af(x))' = a'f(x) + af'(x) = af'(x)$.

And

$$\begin{aligned}\left(\frac{f(x)}{g(x)}\right)' &= \left(f(x) \cdot \frac{1}{g(x)}\right)' = f'(x) \frac{1}{g(x)} + f(x) \left(\frac{1}{g(x)}\right)' \\ &= \frac{f'(x)}{g(x)} + f(x) \frac{-g'(x)}{g(x)^2} = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}.\end{aligned}$$

.

EXERCISES

QUESTION 2

Calculate the derivative of the following functions:

1. $f(x) = \frac{x}{x+1}$

2. $f(x) = \frac{1}{x^2+1}$.

EXP AND LOG

We have $\exp'(x) = \exp(x)$:

$$\begin{aligned}\exp'(x) &= \lim_{h \rightarrow 0} \frac{\exp(x+h) - \exp(x)}{h} = \lim_{h \rightarrow 0} \frac{\exp(x) \exp(h) - \exp(x)}{h} \\ &= \lim_{h \rightarrow 0} \exp(x) \frac{\exp(h) - 1}{h} = \exp(x) \underbrace{\lim_{h \rightarrow 0} \frac{\exp(h) - 1}{h}}_{=1} = \exp(x).\end{aligned}$$

The fact that $\lim_{h \rightarrow 0} \frac{\exp(h)-1}{h} = 1$ follows from the definition of e but is not obvious.

\log is differentiable, although I won't prove it now. Let's assume it's true.

Notice that $\exp(\log(x)) = x$.

Differentiating both sides we get $\exp'(\log(x)) \log'(x) = 1$.

Using $\exp' = \exp$, this is $\exp(\log(x)) \log'(x) = 1$, i.e., $\log'(x) = \frac{1}{x}$.

GENERAL POWER RULE

If $k \in \mathbb{R}$ and $f(x) = x^k$ for $x > 0$ we have

$$\begin{aligned} f'(x) &= (x^k)' \\ &= (\exp(\log(x^k)))' \\ &= (\exp(k \log(x)))' \\ &= \exp'(k \log(x)) k \log'(x) \\ &= \exp(\log(x^k)) k x^{-1} \\ &= x^k k x^{-1} = k x^{k-1}. \end{aligned}$$

QUESTION 3

Calculate the derivative of $f(x) = a^x$ for every $a > 0$.

EXERCISES

QUESTION 4

Calculate the derivative of the following functions:

1. $f(x) = 3x^{1/3}$

2. $f(x) = \frac{1}{100}x^{25} - \frac{1}{10}x^{0.25}$

3. $f(y) = 1 - 1/y^2$

4. $f(x) = \ln(2\pi x^2)$

5. $f(x) = e^{-x^2}$

HIGHER ORDER DERIVATIVES

If $f : U \rightarrow \mathbb{R}$ is **differentiable** (i.e., differentiable at every $x \in U$) then $f' : U \rightarrow \mathbb{R}$ is another function, and we can ask if it is itself differentiable.

Example. Take $f : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(x) = \begin{cases} x^2 & \text{if } x \geq 0, \\ -x^2 & \text{if } x < 0. \end{cases}$$

Notice that it is differentiable. This is only non-obvious at 0. We have

$$\begin{aligned} \lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h} &= \lim_{h \rightarrow 0^+} \frac{h^2}{h} = \lim_{h \rightarrow 0^+} h = 0, \quad \text{and} \\ \lim_{h \rightarrow 0^-} \frac{f(0+h) - f(0)}{h} &= \lim_{h \rightarrow 0^-} \frac{-h^2}{h} = \lim_{h \rightarrow 0^-} -h = 0, \end{aligned}$$

$$\text{so } f'(0) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = 0.$$

HIGHER ORDER DERIVATIVES

We obtain

$$f'(x) = \begin{cases} 2x & \text{if } x \geq 0, \\ -2x & \text{if } x < 0. \end{cases}$$

In other words, $f'(x) = 2|x|$. This is not differentiable at 0. So f' is not differentiable.

But if f' is differentiable at x we can define the **second derivative** $f''(x)$ of f as the derivative of f' at x .

QUESTION 5

If $f(x) = x^3$ calculate $f''(x)$ for all $x \in \mathbb{R}$.

OPTIMIZATION

If $f : D \rightarrow \mathbb{R}$ is maximized or minimized at x in the interior of D and $f'(x)$ exists then $f'(x) = 0$.

The reason is that if $f'(x) > 0$ then when $h > 0$ is sufficiently small we have $\frac{1}{h}(f(x+h) - f(x)) \approx f'(x) > 0$, so $f(x+h) > f(x)$, and $f(x)$ is not the maximum value of f . If $f'(x) < 0$ then when $h < 0$ is small we have $\frac{1}{h}(f(x+h) - f(x)) \approx f'(x) < 0$, so multiplying by h we get $f(x+h) - f(x) > 0$, and again $f(x)$ is not the maximum.

If $f : (a, b) \rightarrow \mathbb{R}$ is differentiable and $f' > 0$ then f is increasing. (Intuition: $f'(x) > 0$ means that f has a “positive slope” at x .)

If $f' < 0$ then f is decreasing.

This is very useful! Example: what is the maximum of $f(x) = x - x^2$? We have $f'(x) = 1 - 2x$, so $f' > 0$ if $x < \frac{1}{2}$ and $f' < 0$ if $x > \frac{1}{2}$. Hence f increases and then decreases. Therefore the max is at $x = \frac{1}{2}$.


EXAMPLE

QUESTION 6

Find the minimum and maximum of $f : [0, 1] \rightarrow \mathbb{R}$ given by $f(x) = 2x^2 - x + 1$.

CONVEXITY


If f is twice differentiable then we say it's **convex** if $f'' \geq 0$, and **concave** if $f'' \leq 0$.



Convex



Concave



Both

If f is concave and $f'(x) = 0$ then f is maximized at x .

If f is convex and $f'(x) = 0$ then f is minimized at x .

If f'' exists around x , is continuous at x , and $f''(x) < 0$ then $f'' < 0$ around x , so f is concave around x , and, so, if $f'(x) = 0$ then f has a **local maximum** at x .

If f'' exists around x , is continuous at x , $f''(x) > 0$, and $f'(x) = 0$ then f has a **local minimum** at x .

EXAMPLE


Take $f(x) = x^3 - 3x + 1$.

We have $f'(x) = 3x^2 - 3$ and $f''(x) = 6x$.

$f'(x) = 0$ is $3x^2 = 3$, i.e., $x = -1$ or $x = 1$.

$f''(-1) = -6 < 0$, so f has a local maximum at -1 .

$f''(1) = 6 > 0$, so f has a local minimum at 1 .




WHAT IF $f''(x) = 0$?

Take $f(x) = x^3$.

We have $f'(x) = 3x^2$ and $f''(x) = 6x$.


We have $f'(0) = 0$ but $f''(0) = 0$, so it's not obvious if it's a local minimum or maximum. It's neither.



Take $f(x) = x^4$.

We have $f'(x) = 4x^3$ and $f''(x) = 12x^2 \geq 0$, so it's convex.

We have $f'(0) = 0$ but $f''(0) = 0$. But f is convex, so it must be minimized at 0.



EXERCISE

QUESTION 7

Find the minimum and maximum of the following functions:

1. $f : [0, 1] \rightarrow \mathbb{R}, f(x) = x(1 - x).$
2. $f : [0, 1] \rightarrow \mathbb{R}, f(x) = x - \sqrt{x}.$
3. $f : [0, 6] \rightarrow \mathbb{R}, f(x) = x^3 - \frac{15}{2}x^2 + 12x + 8.$
4. $f : [-4, 4] \rightarrow \mathbb{R}, f(x) = 3x^4 - 4x^3 - 36x^2.$

L'HÔPITAL'S RULE

THEOREM

Suppose $\lim_{x \rightarrow x_0} f(x) = \lim_{x \rightarrow x_0} g(x)$, both are 0 or ∞ , and f, g are differentiable near x_0 .

Then

$$\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = \lim_{x \rightarrow x_0} \frac{f'(x)}{g'(x)},$$

provided the latter limit exists.

Examples.

$$1. \lim_{x \rightarrow 2} \frac{x^2 - 4}{x^3 - 8} = \lim_{x \rightarrow 2} \frac{2x}{3x^2} = \frac{1}{3}.$$

$$2. \lim_{x \rightarrow +\infty} \frac{e^x}{x^2} = \lim_{x \rightarrow +\infty} \frac{e^x}{2x} = \lim_{x \rightarrow +\infty} \frac{e^x}{2} = +\infty.$$

$$3. \lim_{x \rightarrow 0} x \log(x) = \lim_{x \rightarrow 0} \frac{\log(x)}{1/x} = \lim_{x \rightarrow 0} \frac{1/x}{-1/x^2} = \lim_{x \rightarrow 0} (-x) = 0.$$

EXERCISE

QUESTION 9

Calculate the following limits:

1. $\lim_{x \rightarrow +\infty} \frac{\log(x)}{x}.$

2. $\lim_{x \rightarrow 1} \frac{\sqrt{x} - 1}{x - 1}.$

3. $\lim_{x \rightarrow 0^+} x^x.$

QUESTION 10

Find the minimum and maximum value of $f : (0, 1] \rightarrow \mathbb{R}$ given by $f(x) = x \log(x)$.

